



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore P = \frac{\pi a^2 bc^2 l m n \sin B}{2 \sqrt{[(abl m + ac l n + bcm n)^3]}} = \frac{\pi abc \Delta l m n}{\sqrt{[(abl m + ac l n + bcm n)^3]}}.$$

But $al + bm + cn = 2\Delta$.

$$\therefore P = \frac{\pi \Delta ac l n [2\Delta - al - cn]}{\sqrt{[(2\Delta al + 2\Delta cn - ac l n - a^2 l^2 - c^2 n^2)^3]}}.$$

From the equation to the ellipse, we get

$$BD = u = \frac{acn}{bm + cn} = \frac{acn}{2\Delta - al}, \quad BF = v = \frac{acl}{al + bm} = \frac{acl}{2\Delta - cn}.$$

$$\therefore al = \frac{2\Delta v(a-u)}{ac-uv}, \quad cn = \frac{2\Delta u(c-v)}{ac-uv}.$$

$$\therefore P = \frac{\pi \Delta uv \sqrt{[(a-u)(c-v)]}}{\sqrt{[(av + cu - uv)^3]}}. \quad \text{Let } a-u=t, \quad c-v=z.$$

$$\therefore P = \frac{\pi \Delta [a-t][c-z] \sqrt{[tz]}}{\sqrt{[(ac-tz)^3]}}.$$

$$\therefore Q = \int_0^a \int_0^c P dt dz / \int_0^a \int_0^c dt dz = \frac{\pi \Delta}{ac} \int_0^a \int_0^c \frac{(a-t)(c-z) \sqrt{[tz]} dt dz}{\sqrt{[(ac-tz)^3]}}.$$

Let $tz = ac \sin^2 \theta$, $\theta' = \sin^{-1} \sqrt{(t/a)}$.

$$\begin{aligned} \therefore Q &= \frac{2\pi \Delta}{a} \int_0^a \int_0^{\theta'} \frac{(a-t)(t - a \sin^2 \theta) \sin^2 \theta dt d\theta}{t^2 \cos^2 \theta} \\ &= \frac{\pi \Delta}{a} \int_0^a \left(\frac{a-t}{t^2} \right) \{ 3a \sin^{-1} \sqrt{(t/a)} - 2t \sin^{-1} \sqrt{(t/a)} - 3 \sqrt{[t(a-t)]} \} dt. \end{aligned}$$

Let $t = a \sin^2 \varphi$.

$$\therefore Q = 2\pi \Delta \int_0^{\frac{1}{2}\pi} (3\varphi - 2\varphi \sin^2 \varphi - 3 \sin \varphi \cos \varphi) \cot^3 \varphi d\varphi = \frac{1}{2} \pi^2 \Delta (7 - 10 \log^2)$$

127. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

What is the probable error of the volume of a rectangular parallelepiped whose edges measured by the repeated application of a unit of measure are found to be a, b, c , supposing that the probable error of a line so measured whose length is found to be l is r/l ?

Solution by the PROPOSER.

The probable error for a , $=r_1/a$; for b , r_1/b ; for c , r_1/c .

\therefore Error of volume in length $=bcr_1/a$.

Error of volume in width $=acr_1/b$.

Error of volume in thickness $=abr_1/c$.

The probable error of volume = square root of the sum of the squares of these three errors.

\therefore Probable error $=\sqrt{[(ab^2c^2 + a^2bc^2 + a^2b^2c)r^2]} = r_1/[abc(ab + ac + bc)]$.

MISCELLANEOUS.

124. Proposed by J. W. YOUNG, Graduate Student, Cornell University, Ithaca, N. Y.

Prove that the general value of θ , which satisfies the equation

$$(\cos\theta + i\sin\theta)(\cos 2\theta + i\sin 2\theta) \dots \text{to } n \text{ factors} = 1 \text{ is } \frac{4m\pi}{n(n+1)};$$

where m is any integer ($i = \sqrt{-1}$).

Solution by G. W. GREENWOOD, A. M., McKendree College, Lebanon, Ill.; LON C. WALKER, A. M., Leland Stanford Jr. University, Cal., and J. SCHEFFER, A. M., Hagerstown, Md.

$$\begin{aligned} 1 &= (\cos\theta + i\sin\theta)(\cos 2\theta + i\sin 2\theta) \dots (\cos n\theta + i\sin n\theta) = (\cos\theta + i\sin\theta)^{1+2+\dots+n} \\ &= (\cos\theta + i\sin\theta)^{\frac{1}{2}n(n+1)} = \cos \frac{n(n+1)\theta}{2} + i \sin \frac{n(n+1)\theta}{2}. \end{aligned}$$

$$\therefore \frac{n(n+1)\theta}{2} = 2m\pi, \text{ where } m \text{ is any integer; i. e. } \theta = \frac{4m\pi}{n(n+1)}.$$

Also solved by G. B. M. ZERR.

125. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Assume $m = nt + \varepsilon - \omega$, thus giving $v = m + e \sin v$ as the relation connecting the mean and eccentric anomalies, then express $x = a \cos v$, $y = b \sin v$, and $r = a(1 - e \cos v)$ by a Fourier series in terms of m .

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

If $y_1 = z + x\varphi((y))$, we get by Lagrange's Theorem,

$$\begin{aligned} f(y_1) &= f(z) + x\varphi(z)f'(z) + \frac{x^2}{1.2} \frac{d}{dz} \{[\varphi(z)]^2 f''(z)\} + \frac{x^3}{1.2.3} \left(\frac{d}{dz}\right)^2 \{[\varphi(z)]^3 f''(z)\} + \\ &\quad \text{etc., etc.} \end{aligned}$$

From $v = m + e \sin v$, $y_1 = v$, $z = m$, $x = e$, $\varphi(y) = \sin v$.

Now $f(v) = v$ and $f'(v) = 1$.